Innovation for Our Energy Future

Solar Radiation Measurements:

A Workshop For

The National Association of State Universities and Land Grant Colleges

By

Tom Stoffel & Steve Wilcox

Hydrogen & Electric Technologies & Systems Center August 4, 2004



Outline

- Introductions
- Shining On, A Primer on Solar Radiation Data
 - What are solar radiation measurements?
 - Why do we need solar radiation data?
 - What influences the amount of solar radiation?
 - How do we use solar radiation data?
 - How accurate do the data need to be?
- How are we meeting our solar radiation data needs?
- Where can you obtain solar radiation data?
- Pop Quiz
 - No acronyms!



Introductions

Tom Stoffel & Steve Wilcox

Resource Integration Group

Measurement & Instrumentation Team

Geographic Information System Team

40⁺ years experience:

- Solar measurement station/network design
 - SRRL, HBCU, Saudi, DOE/ARM, NOAA, WMO/BSRN, GAW
- Radiometer calibration and characterization
 - BORCAL/RCC
 - IPCs, NPCs
- Solar data quality assessment
 - SERI-QC
 - DQMS



What are Solar Radiation Measurements?

Energy from the Sun at the Earth's Surface

- Different parts of the sky
- Change with time (minutes, hours)
- Change with time (seasons, years, decades)
- Change with location




What are Solar Radiation Measurements?

Light from the sky dome

- Direct from the sun
- Everywhere but the sun
- Entire sky

We call it

- Direct (beam)
- Diffuse (sky)
- Global (total)



Global is the sum of direct and diffuse



What are Solar Radiation Measurements?

Direct Normal

Measured by a

Pyrheliometer on a

sun-following

tracker

Global Horizontal

Measured by a Pyranometer with a horizontal sensor

Diffuse

Measured by a shaded *Pyranometer* under a tracking ball









Solar Irradiance Components

Global = Direct Normal * Cos(Z) + Diffuse





Cos(Z) = 8/18


 $Z = Cos^{-1}(0.4444)$


 $Z = 63.6^{\circ}$

18 dots

8 dots

Clear Sky






http://www.nrel.gov/srrl

Thermopile Detectors

How do the radiometers work? Thermo-electric detectors:

Two metals + Heat = Electrical Current

Copper-Constantan wire wound *Thermopiles*




Thermopile Detectors

Pyrheliometer

1st Class \$, Flat Spectral Response, "Slow"

Pyranometer



The Eppley Laboratory, Inc.

Photoelectric Detectors



www.kippzonen.com



www.licor.com

Fast, Low-Cost, with Reduced Spectral Response:






Figure 4. The LI-200SA Pyranometer spectral response is illustrated along with the energy distribution in the solar spectrum (8).

Partly Cloudy Sky




http://www.nrel.gov/srrl


Changes with Time & Location: Annual Cycle



Changes with Time: Inter-annual



Changes with Time: Inter-annual



Spectral Distribution of Solar Radiation

Broadband Solar Radiation:

280 nm - 3,000 nm




TIFF (Uncompressed) decompressor are needed to see this picture.

(99% of "shortwave" irradiance at the surface)


Spectral Irradiance

Basic Solar Spectral Regions:


- Ultraviolet.....200 400 nm
- Visible......400 700 nm
- Infrared......700 3000 nm




Follow the Photons!




Photovoltaic Responses



Simple Model for Atmospheric Radiative Transfer of Sunshine SMARTS



SMARTS



Available from NREL: http://rredc.nrel.gov

Why Do We Need Solar Radiation Data?

Agriculture

Astronomy

Atmospheric Science

Climate Change

Health

Hydrology

Materials

Oceanography

Photobiology

Renewable Energy

Photosynthesis

Solar Output Variation

Numerical Weather Prediction

Energy Balance

UV effects on skin

Evaporation

Degradation

Energy Balance

Light and Life


Sustainability

Why Do We Need Solar Radiation Data? Renewable Energy

The amount of solar energy reaching the earth's land areas in 1 hour is enough to supply the U.S. energy needs for

1 year (~100 Quads/yr)

- Photovoltaics
- Solar Heat-thermal
- Solar Heat-electric
- Solar Fuel-biomass
- Passive Solar Lighting
- Building HVAC
- Solar Detoxification



What Influences the Amount of Solar Radiation?

- Solar output
- Earth-Sun distance
- Clouds
- Water vapor
- Air pollution
- Smoke from forest fires
- Volcanic ash
- Location
- Time of day
- Season

11 year solar cycle

3.5% annual variation

Dominant factor

Selective absorber


40% less direct

Natural or man-made

Global effect for years

Solar position


Solar "Constant"



World Radiation Center, Davos, Switzerland http://www.pmodwrc.ch/




What Influences the Amount of Solar Radiation?




Earth's Orbit:

- Earth-Sun distance
- Relative tilt
- Time of day



What Influences the Amount of Solar Radiation?



How Do We Use Solar Radiation Data?

- Technology Selection
- Siting
- System Design
- Performance Monitoring











PV Energy kWh/kW-yr

Effective Load Carrying Capacity







Source: Christy Herig (NREL) and Richard Perez (SUNY/Albany)

- PV can provide peak shaving in many parts of U.S.
- During off-peak periods, PV capacity can be applied to hydrogen generation






An Integrated Analysis Utilizing GIS can Assist With Energy and Environment Planning Efforts




Satellite-Derived Techniques Provide Improved Site-Time Coverage (SUNY/Albany)




GLOBAL IRRADIANCE (average W/sq.m)

DIRECT IRRADIANCE (average W/sq.m)



How Do We Use Solar Radiation Data?

General Circulation Model Development



DOE/Atmospheric Radiation Measurement (ARM) Program



How Accurate Do the Data Need to Be?

- What are the risks?
 - Cost/Benefit of Resource Assessment approach
- What is the application?
 - Daylighting & building thermal performance
 - Concentrating Collector Solar Power Plant
 - Cloud forcing analyses for climate change research
- What is the period of interest?
 - Measurement uncertainties decrease with longer averaging intervals (averaging can remove random errors)
 - Recent data more accurate than historical records (technology advancements)



How Accurate Do the Data Need to Be?

What is possible?

Measurement Uncertainty Estimates*

	Pyrheliometer (Direct Normal)	Pyranometer (Global)
Calibration	±1.6%	±4.2%
Field Data (Best practice)	~ ±5%	~ ±5%

Instantaneous data intervals

How Will We Meet Our Solar Radiation Data Needs?

Research Activities:

- Solar Radiation Research Laboratory
 - Metrology
 - Optics
 - Electronics
 - Data Acquisition
- Photovoltaic Program
 - Radiometric Measurements
- Climate Change
 - Broadband Radiometer Mentor
- Collaborations
 - WMO, UNEP, NCAR, NOAA, state & local govt, academia



Solar Radiation Research Lab



- Baseline Measurements
- Radiometer Calibrations
- Instrument Development
- Station Operator Training



Solar Radiation Research Lab



Baseline Measurements
(98 data elements)
http://www.nrel.gov/midc



Rotating Shadowband Pyranometer

Radiometer Calibrations



World Radiometric Reference




NREL Transfer Standards

NPC At SRRL




Radiometer Calibrations



National Solar Radiation Data Base




NSRDB Stations (1961-1990)



Solar Measurement Stations (1990 - Present)

Automatic Data Quality



Where Can You Obtain solar Radiation Data?

- Renewable Resource Data Center
 - http://rredc.nrel.gov
- Measurement & Instrumentation Data Center
 - http://www.nrel.gov/midc
- NREL Map Server
 - http://www.nrel.gov/maps
- World Radiation Data Center
 - http://wrdc-mgo.nrel.gov
- National Climatic Data Center
 - http://www.ncdc.noaa.gov
- DOE Atmospheric Radiation Measurement Program
 - http://www.arm.gov
- NOAA Climate Monitoring & Diagnostic Laboratory
 - http://www.cmdl.noaa/gov/star
- NOAA Surface Radiation Research Branch
 - http://www.srrb.noaa.gov

Key Points

- Accurate information is important for policy decisions, technology selection, siting, designing, and monitoring the performance of solar energy conversion systems
- Accurate measurements are important for model development
- The work we do to improve solar measurements
 - Calibration
 - Instrument characterization
 - Measurement techniques (operations and maintenance, radiometer selection, installation considerations, etc.)
 - Data Quality Assessment
 - Training
- Data distribution to meet user needs (MIDC, RReDC, NSRDB)

Solar Radiation Measurement

Thank you!

Questions?


Write the relationship between Global, Direct, & Diffuse irradiance.

Global = Direct Normal * Cos(Z) + Diffuse

Indicate which properties (quick, accurate, or cheap) apply to these pyranometer detector types:

- ✓ Photodiode Fast, Cheap, Spectrally selective
- ✓ Thermopile Accurate, \$\$, Slow

T/F: The Global irradiance can never exceed the solar constant.



The presently accepted value of the Solar Constant:

- a) 1.96 Langleys per minute
- b) 1366 Watts per square meter
- c) 432.7 BTUs per hour-square foot
- d) All of the above

